

Lake Minnetonka Pet Wellness Center Wayzata Phone: 952 471-0911 www.tonka.vet Westonka Animal Hospital & Laser Surgery Center Mound Phone: 952 472-4900 www.westonka.vet



Anal Gland Disease In Pets

What Are Anal Sacs?

The anal sacs are located on either side of the anus at the 8 o'clock and 4 o'clock positions; they are positioned just under the skin. They connect to the anus by means of small canals or ducts. Anal sacs produce and store a dark, foul-smelling fluid. These are the same type of organs that a skunk has to scare away its enemies. Although pets can use these for the same purpose, most pets live in an environment that has no enemies. Because the sacs are rarely emptied, the fluid builds up, solidifies, and becomes an ideal environment in which bacteria can grow.

What disorders can occur in the anal sacs?

There are three diseases that occur in the anal sacs.

- 1) When the fluid becomes thick and solidified, the condition is called **impaction**.
- 2) When bacteria grow in this material producing a yellow or bloody pus, the condition is called **infection**.
- 3) When the infection builds to create a hot, tender swelling in the gland, the condition is called an **abscess**. When the abscessed material overflows the sac, the skin over the sac breaks open, and the pus drains onto the skin.

How will I know if my pet is having problems with its anal sacs?

Symptoms of anal sac disease are:

- 1) Scooting or dragging the anal area.
- 2) Excessive licking under the tail.
- 3) Pain, sometimes severe, near the tail or anus.
- 4) A swollen area on either side of the anus.
- 5) Bloody or sticky drainage on either side of the anus.

Enlarged, Inflamed Anal Sac Abscessed Anal Sac

How are the various anal sac diseases treated?

The treatment for **impaction** is to express the sacs and clean out the solidified material. For **infection**, the sacs must be expressed and antibiotics administered to kill the bacteria. If the sacs **abscess**, the abscess must be surgically drained and antibiotics administered.

My pet has had several bouts of anal sac disease. Is there a long-term cure?

Many pets have recurrent anal sac disease. Some breeds of dogs, such as Poodles, commonly have problems. The anal sacs of obese pets do not drain well, and thus these pets are predisposed to recurrent problems. If a pet has several episodes of anal sac disease, the anal sacs can be removed surgically. Because these sacs are virtually unused, there is no loss to the dog or cat. It is the only way to permanently cure the problem.

Are there any complications with the surgery?

Surgery requires general anesthesia which always carries some degree of risk, whether the patient is a pet or a person. However, modern anesthetics make this risk very minimal for pets that are otherwise healthy.

Some pets will experience temporary lack of good bowel control. They may drop fecal balls as they walk. This occurs because the nerves that control the anus are very near the anal sacs and may be damaged during surgery. However, this is almost always a temporary problem that will resolve itself in a few days to a couple of weeks.

My pet frequently leaves a foul-smelling drop of liquid on the furniture. Is this related?

Some dog and cats are born with anal canals that do not close well. These pets are constantly draining anal sac fluid and leaving a foul-smelling drop wherever they have been. This is another indication for anal sac removal since there does not appear to be any other way to stop this, and these pets do not outgrow this problem.